


This listing of claims will replace all prior versions, and listings, of claims in the application:

**LISTING OF CLAIMS:**

1. (Previously presented) A multireactive polymerizable mesogenic compound of formula I



wherein

$R^1$  is halogen, CN, OCN, NCS,  $NO_2$  or an alkyl radical with 1 to 30 C atoms which may be unsubstituted, mono- or polysubstituted by halogen or CN, optionally one or more non-adjacent  $CH_2$  groups being replaced, in each case independently from one another, by -O-, -S-, -NH-, -N( $CH_3$ )-, -CO-, -COO-, -OCO-, -OCO-O-, -S-CO-, -CO-S-, -CH=CH- or -C $\equiv$ C- in such a manner that oxygen atoms are not linked directly to one another, or alternatively has one of the meanings of  $R^2$  or is P-(Sp-X) $_n$ ,

P is a polymerizable group,

Sp is a spacer group with 1 to 25 C atoms,

X is -O-, -S-, -CO-, -COO-, -OCO-, -OCO-O-, -CO-NH-,  
-NH-CO-, -OCH $_2$ -, -CH $_2$ O-, -SCH $_2$ -, -CH $_2$ S-, -CH=CH-COO-, -OOC-CH=CH-  
or a single bond,

n is 0 or 1,

MG is a mesogenic group, and

$R^2$  is straight-chain or branched alkyl with 1 to 25 C atoms which may be unsubstituted, mono- or polysubstituted by halogen or CN, optionally one or more non-adjacent  $CH_2$  groups being replaced, in each case independently from one another, by -O-,

-S-, -NH-, -N(CH<sub>3</sub>)-, -CO-, -COO-, -OCO-, -OCO-O-, -S-CO-, -CO-S-, -CH=CH- or -C≡C- in such a manner that oxygen atoms are not linked directly to one another, and which is substituted with at least two identical or different groups P.

**2. (Previously presented)** A multireactive polymerizable mesogenic compound according to claim 1, wherein R<sup>1</sup> is not a polymerizable group.

**3. (Original)** A multireactive polymerizable mesogenic compound according to claim 1, wherein R<sup>1</sup> has one of the meanings of R<sup>2</sup>.

**4. (Original)** A multireactive polymerizable mesogenic compound according to claim 1, wherein MG is of formula II



wherein

Z is -O-, -S-, -CO-, -COO-, -OCO-, -CO-NH-, -NH-CO-,  
-CH<sub>2</sub>CH<sub>2</sub>-, -OCH<sub>2</sub>-, -CH<sub>2</sub>O-, -SCH<sub>2</sub>-, -CH<sub>2</sub>S-, -CH=CH-,  
-CH=CH-COO-, -OCO-CH=CH-, -C≡C- or a single bond,

A<sup>1</sup> and A<sup>2</sup> are each independently 1,4-phenylene in which, in addition, one or more CH groups are optionally replaced by N; 1,4-cyclohexylene in which, in addition, one or two non-adjacent CH<sub>2</sub> groups are optionally replaced by O and/or S; 1,4-cyclohexenylene; 1,4-bicyclo(2,2,2)octylene; piperidine-1,4-diyl; naphthalene-2,6-diyl; decahydronaphthalene-2,6-diyl; or 1,2,3,4-tetrahydro-naphthalene-2,6-diyl; all these groups optionally being

unsubstituted, mono- or polysubstituted with F, Cl, OH, CN, NO<sub>2</sub> or alkyl, alkoxy, alkylcarbonyl or alkoxy carbonyl groups having 1 to 7 C atoms wherein one or more H atoms may be substituted by F or Cl, and

m is 1, 2 or 3.


5. **(Original)** A multireactive polymerizable mesogenic compound according to claim 1, wherein P is selected from CH<sub>2</sub>=CW-COO-, WCH=CH-O-, CH<sub>2</sub>=CH-Phenyl-(O)<sub>k</sub>-

and  $\text{WHC} \begin{array}{c} \diagup \text{O} \diagdown \\ \text{---} \end{array} \text{CH=}$ , with W being H, CH<sub>3</sub> or Cl and k being 0 or 1.

6. **(Original)** A multireactive polymerizable mesogenic compound according to claim 1, wherein R<sup>2</sup> is substituted with 2, 3, 4 or 5 identical or different polymerizable groups P.

7. **(Previously presented)** A multireactive polymerizable mesogenic compound according to claim 1, wherein R<sup>2</sup> is a group of one of the following formulae

-X-alkyl-CHP <sup>1</sup> -CH <sub>2</sub> -CH <sub>2</sub> P <sup>2</sup>	Ia
-X-alkyl-C(CH <sub>2</sub> P <sup>1</sup> )(CH <sub>2</sub> P <sup>2</sup> )-CH <sub>2</sub> P <sup>3</sup>	Ib
-X-alkyl-CHP <sup>1</sup> CHP <sup>2</sup> -CH <sub>2</sub> P <sup>3</sup>	Ic
-X-alkyl-C(CH <sub>2</sub> P <sup>1</sup> )(CH <sub>2</sub> P <sup>2</sup> )-C <sub>a</sub> H <sub>2a+1</sub>	Id
-X-alkyl-CHP <sup>1</sup> -CH <sub>2</sub> P <sup>2</sup>	Ie
-X-alkyl-CHP <sup>1</sup> P <sup>2</sup>	If
-X-alkyl-CP <sup>1</sup> P <sup>2</sup> -C <sub>a</sub> H <sub>2a+1</sub>	Ig
-X-alkyl-C(CH <sub>2</sub> P <sup>1</sup> )(CH <sub>2</sub> P <sup>2</sup> )-CH <sub>2</sub> OCH <sub>2</sub> -C(CH <sub>2</sub> P <sup>3</sup> )(CH <sub>2</sub> P <sup>4</sup> )CH <sub>2</sub> P <sup>5</sup>	Ih
-X-alkyl-CH((CH <sub>2</sub> ) <sub>a</sub> P <sup>1</sup> )((CH <sub>2</sub> ) <sub>b</sub> P <sup>2</sup> )	Ii



Ik

wherein



alkyl is straight-chain or branched alkylene with 1 to 12 C atoms which may be unsubstituted, mono- or polysubstituted by halogen or CN, one or more non-adjacent CH<sub>2</sub> groups optionally being replaced, in each case independently from one another, by -O-, -S-, -NH-, -N(CH<sub>3</sub>)-, -CO-, -COO-, -OCO-, -OCO-O-, -S-CO-, -CO-S-, -CH=CH- or -C≡C- in such a manner that oxygen atoms are not linked directly to one another,

a and b are identical or different integers from 0 to 6,

X has one of the meanings given in formula I, and

P<sup>1</sup> to P<sup>5</sup> independently have one of the meanings of P given in formula I.

**8. (Previously presented)** A multireactive polymerizable mesogenic compound according to claim 5, wherein R<sup>2</sup> is a group of one of the following formulae





Ik

wherein

alkyl is straight-chain or branched alkylene with 1 to 12 C atoms which may be unsubstituted, mono- or polysubstituted by halogen or CN, one or more non-adjacent CH<sub>2</sub> groups optionally being replaced, in each case independently from one another, by -O-, -S-, -NH-, -N(CH<sub>3</sub>)-, -CO-, -COO-, -OCO-, -OCO-O-, -S-CO-, -CO-S-, -CH=CH- or -C≡C- in such a manner that oxygen atoms are not linked directly to one another,

a and b are identical or different integers from 0 to 6,

X has one of the meanings given in formula I, and

P<sup>1</sup> to P<sup>5</sup> independently have one of the meanings of P given in formula I.

**9. (Previously presented)** A multireactive polymerizable mesogenic compound according to claim 7, wherein alkyl is -(CH<sub>2</sub>)<sub>c</sub>-, with c being an integer from 1 to 12.

**10. (Original)** A multireactive polymerizable mesogenic compound according to claim 1, wherein each P is independently of each other acrylate, methacrylate, vinyl, vinyloxy, epoxy or p-vinylphenyloxy.

**11. – 16. (CANCELED)**

**17. (Previously presented)** A multireactive polymerizable mesogenic compound according to claim 8, wherein alkyl is -(CH<sub>2</sub>)<sub>c</sub>-, with c being an integer from 1 to 12.

**18. (Previously presented)** A multireactive polymerizable mesogenic compound according to claim 1, wherein MG is a group of one of the following formulae II-1 to II-25 or a mirror image thereof:

-Phe-Z-Phe-	II-1
-Phe-Z-Cyc-	II-2
-Cyc-Z-Cyc-	II-3
-PheL-Z-Phe-	II-4
-PheL-Z-Cyc-	II-5
-PheL-Z-PheL-	II-6
-Phe-Z-Phe-Z-Phe-	II-7
-Phe-Z-Phe-Z-Cyc-	II-8
-Phe-Z-Cyc-Z-Phe-	II-9
-Cyc-Z-Phe-Z-Cyc-	II-10
-Phe-Z-Cyc-Z-Cyc-	II-11
-Cyc-Z-Cyc-Z-Cyc-	II-12
-Phe-Z-Phe-Z-PheL-	II-13
-Phe-Z-PheL-Z-Phe-	II-14
-PheL-Z-Phe-Z-Phe-	II-15
-PheL-Z-Phe-Z-PheL-	II-16
-PheL-Z-PheL-Z-Phe-	II-17
-PheL-Z-PheL-Z-PheL-	II-18
-Phe-Z-PheL-Z-Cyc-	II-19
-Phe-Z-Cyc-Z-PheL-	II-20
-Cyc-Z-Phe-Z-PheL-	II-21
-PheL-Z-Cyc-Z-PheL-	II-22
-PheL-Z-PheL-Z-Cyc-	II-23
-PheL-Z-Cyc-Z-Cyc-	II-24
-Cyc-Z-PheL-Z-Cyc-	II-25

wherein Phe is 1,4-phenylene, PheL is a 1,4-phenylene group which is substituted by 1 to 4 groups L, with L being F, Cl, CN, OH, NO<sub>2</sub> or an optionally fluorinated alkyl, alkoxy or alkanoyl group with 1 to 7 C atoms, Cyc is 1,4-cyclohexylene and Z are independently -O-,

-S-, -CO-, -COO-, -OCO-, -CO-NH-, -NH-CO-, -CH<sub>2</sub>CH<sub>2</sub>-, -OCH<sub>2</sub>-, -CH<sub>2</sub>O-, -SCH<sub>2</sub>-,  
-CH<sub>2</sub>S-, -CH=CH-, -CH=CH-COO-, -OCO-CH=CH-, -C≡C- or a single bond.

**19. (Previously presented)** A multireactive polymerizable mesogenic compound according to claim 1, wherein Sp is a linear or branched alkylene group having 1 to 20 C atoms, in which one or more non-adjacent CH<sub>2</sub> groups are optionally replaced by -O-, -S-, -NH-, -N(CH<sub>3</sub>)-, -CO-, -O-CO-, -S-CO-, -O-COO-, -CO-S-, -CO-O-, -CH(halogen)-, -CH(CN)-, -CH=CH- or -C≡C-.

**20. (Previously presented)** A multireactive polymerizable mesogenic compound according to claim 1, wherein R<sup>1</sup> is a chiral alkyl radical with 1 to 30 C atoms which may be unsubstituted, mono- or polysubstituted by halogen or CN, optionally one or more non-adjacent CH<sub>2</sub> groups being replaced, in each case independently from one another, by -O-, -S-, -NH-, -N(CH<sub>3</sub>)-, -CO-, -COO-, -OCO-, -OCO-O-, -S-CO-, -CO-S-, -CH=CH- or -C≡C- in such a manner that oxygen atoms are not linked directly to one another.